

WO9920795

Publication Title:

METHODS FOR DETERMINING COAT COLOUR GENOTYPES IN PIGS

Abstract:

Abstract of WO9920795

Methods for determining coat colour genotypes in pigs are provided. In particular, these methods are based on determining whether a mutation is/is not present at one or more exon/intron splice sites of the KIT gene. Kits for carrying out such methods are also described. Data supplied from the esp@cenet database - Worldwide

Courtesy of <http://v3.espacenet.com>

This Patent PDF Generated by Patent Fetcher(TM), a service of Stroke of Color, Inc.

ATTORNEY DOCKET NUMBER: 10589-013-999
SERIAL NUMBER: 10/551,301
REFERENCE: **B03**

BEST AVAILABLE COPY

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, G01N 33/68	A1	(11) International Publication Number: WO 99/20795 (43) International Publication Date: 29 April 1999 (29.04.99)
(21) International Application Number: PCT/GB98/03081 (22) International Filing Date: 13 October 1998 (13.10.98) (30) Priority Data: 9722027.1 17 October 1997 (17.10.97) GB (71) Applicant (for all designated States except US): MELICA HB [SE/SE]; Bergagatan 30, S-752 39 Uppsala (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): ANDERSSON, Leif [SE/SE]; Melica HB, Bergagatan 30, S-752 39 Uppsala (SE). MARKLUND, Stefan [SE/US]; 106-D University Village, Ames, IA 50010 (US). KIJAS, James [AU/SE]; Borjegatan 54B, S-752 29 Uppsala (SE). MOLLER, Maria [SE/SE]; Karlsrogatan 46, S-752 39 Uppsala (SE). WALES, Richard [GB/GB]; Pic International Group, c/o Du Pont (U.K.) Ltd., Cereals Innovation Centre, Block B, The Mill Site, 40 Station Road, Cambridge CB1 2JN (GB). (74) Agents: CHAPMAN, Paul, William et al.; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHODS FOR DETERMINING COAT COLOUR GENOTYPES IN PIGS (57) Abstract Methods for determining coat colour genotypes in pigs are provided. In particular, these methods are based on determining whether a mutation is/is not present at one or more exon/intron splice sites of the KIT gene. Kits for carrying out such methods are also described.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS FOR DETERMINING COAT COLOUR GENOTYPES IN PIGS

The present invention relates to methods for determining coat colour genotype in pigs. In particular, it relates to methods of distinguishing between the alleles
5 *I*, *P*, *I*^{*} and *i* of the *KIT* gene.

Coat colour is important to the pig breeding industry for a number of reasons. Firstly in a number of markets there is a preference for white skinned meat. This is due to the fact that pork is often marketed with the skin still attached,
10 and skins from coloured pigs, even if dehaired, can still exhibit coloured hair roots, which can lead to negative perception by the consumer, since the surface of the meat may appear to be spotted by mould. It is therefore necessary in these markets to remove the skin from such carcasses, entailing additional cost.

15 For example, in the US, coloured carcasses are associated with approximately 1% of skin defects requiring dehairing and skinning to remove pigment. As a result of this, coloured pig carcasses are generally discounted. Secondly gross variation in the appearance of pigs claimed to be genetically consistent for other traits can lead to questions about the consistency and quality of the animals in
20 the mind of pig-producing customers. Breeders would also like to be able to ensure consistency in breeding populations. Thus breeders may wish to ensure that progeny produced by breeding crosses were always white. Alternatively a breeder producing a coloured breed may wish to ensure that the correct coat colour characteristics were maintained even during the introgression of genes
25 from white lines.

White coat colour in pigs is controlled by the dominant white locus designated *I* (for inhibition of coat colour). Structural alterations in the porcine *KIT* gene have recently been correlated with various alleles of *I* and are probably responsible for the differences in coat colour pattern found or are closely linked to the mutations that are found (Johansson Moller *et al.* 1996 Mammalian Genome 7, 822-830). Four structural versions of the porcine *KIT* gene have been identified to date and designated *I*, *P*, *I*^{*} and *i* (see figure 1 and table 1). The version found in fully coloured animals including wild boar and which is therefore generally accepted as the wild type allele is *i*. The other versions of the gene known all involve duplication of at least part of the porcine *KIT* gene. *P* is a partially dominant allele and causes in the heterozygous state (*P/i*) a phenotype "patch" characterised by patches of white and coloured coat. *I* and *I*^{*} are both fully dominant and cause a white phenotype both in the heterozygous and homozygous state. The only difference between *I* and *I*^{*} is that there is a 4bp deletion in intron 18 in one of the two *KIT* gene copies associated with the *I* allele. This sequence polymorphism is not expected to have any functional effect.

The phenotypes of a number of gene combinations are listed below with the *I* allele being dominant over *P* and *i*, and *P* being dominant over *i* (Johansson *et al.* 1992 Genomics 14, 965-969)

	<u>Genotype</u>	<u>Colour</u>
	<i>I/I</i>	White
	<i>I/P</i>	White
25	<i>I/i</i>	White

<i>P/i</i>	Patched
<i>i/i</i>	Coloured
<i>I*/i</i>	White

5 Previously Moller Johansson et al. (1996 Mammalian Genome 7, 822-830)
revealed that KIT occurs as a single copy gene in coloured (*i/i*) animals but is
duplicated in the *I*, *I** and *P* allele. This duplication allowed the differentiation
of *I*, *I** and *P* from *i* through examination of the gene copy number of KIT and
to use linked polymorphisms in or in the near vicinity of the KIT gene to
10 distinguish different alleles at the *I* locus. This approach is the subject of
International patent WO97/05278.

A problem remaining from this method however is that it does not distinguish
between *I** and *P*. The result of this is that if one wishes to use the screen to
15 remove heterozygous carrier animals of genotype *I/P* from a white pig line one
also excludes *I/I** animals unnecessarily. The removal of animals, potentially
very valuable if at the top of a breeding pyramid, unnecessarily can lead to a
reduction in the rate of improvement of other traits within that particular group
of animals. Other consequences include a general loss of genetic diversity and a
20 loss of alleles at the locus in question that may have as yet undetected value. It
is essential that one only excludes alleles where absolutely necessary.

Further sequence analysis of genomic DNA revealed a mutation at the first
nucleotide in intron 17 of KIT2 leading to a defect in the splicing of RNA
25 transcribed from this particular copy of the KIT gene. In experiments involving

several pig breeds, this mutation showed a complete concordance with the presence of the *I* and *I** alleles but was not found in the *i* and *I*^p alleles

5 The majority of eukaryotic genes are composed of both coding (exon) and intervening noncoding (intron) regions. The latter of these sequences are removed from large pre-mRNA by a highly accurate cleavage and ligation reaction known as splicing. The result is a mature mRNA transcript, devoid of intron sequences, which is transported to the cytoplasm for translation. The splicing of eukaryotic genes is most likely a two step procedure. First, the pre-mRNA is cleaved at the 5' (donor) splice site followed by cleavage at the 3' (acceptor) splice site. The second step involves rejoining of the spliced exons to result in exclusion of the intervening intron. Splicing is therefore critically dependant on the accuracy of the cleavage and ligation reactions. This accuracy appears to be dependant on the almost completely invariant GT and AG dinucleotides present at the 5' and 3' exon/intron boundaries respectively. These dinucleotides and the often highly conserved surrounding sequences are known as splice sites and serve to bind the protein factors required to perform the cleavage and ligation reactions.

10 15 20 25 The consequences for mutation occurring within a splice site may be a reduction in the amount of mature mRNA produced and/or utilization of alternative but incorrect splice sites in the vicinity. The result is production of mRNA which either contains additional intron sequence or which may lack a portion of coding sequence. Where mutation occurs within the 5' (donor) splice site and prevents binding of protein factors, the exon is no longer recognised as such and is excised along with its neighbouring introns. This is referred to as

'exon skipping' (as reviewed by Cooper and Krawczak, 1994 Human Gene Mutation. Bios Scientific Publishers, Oxford, UK, 1994).

5 In the mutation identified here the G of the conserved GT pair at the exon17/intron17 boundary region is altered to an A. That there is an alteration in the messenger RNA that is translated into protein and that the presence of the change correlates with the white/patched phenotypes suggests that this is the functional mutation rather than merely a linked marker. The splice variant is expected to give rise to a defective protein as 41 amino acids in the mature
10 protein are missing. We therefore assume that this splice mutation is the causal mutation for the difference between *I* and *P*. Our current knowledge as regards molecular differences between alleles at the *I* locus is summarised in Table 1. In conclusion, we have identified two functionally important mutations. One is the gene duplication present in *P*, *I*^{*} and *I* which by itself appears to cause the
15 patch phenotype. The exact reason for this phenotypic effect is not known but one can speculate on the basis of comparative data from the mouse that the duplicated copy of the KIT gene may lead to a defect in gene expression which in turn affects melanocyte migration. This is especially valuable in that there can be no breakdown of the linkage between the DNA polymorphism and the
20 trait itself. This has allowed us to develop a range of assays for the determination of the presence of the DNA polymorphism and the genotype of the animal with regard to coat colour determination to a significantly greater extent than has been possible before. The second mutation is the splice mutation that occurs in one of the KIT copies associated with the dominant white (*I* and
25 *I*^{*}) alleles. The expression of a truncated form of the KIT protein is expected to cause a more severe defect in KIT function and a more severe effect on coat colour.

Table 1.

	Allele	KIT gene	Intron 17	Associated Phenotype
5	<i>i</i>	KIT1	Normal	Coloured
	<i>P</i>	KIT1 KIT2	Normal Normal	Patch
10	<i>I</i>	KIT1 KIT2	Normal Mutated	Dominant White
15	<i>I*</i>	KIT1 KIT2	Normal Mutated	Dominant White

In addition to these two functionally important mutations, a mutation in the KIT gene with no known phenotypic effect, the 4 bp deletion in intron 18 has been documented at the *I* locus as described in International patent application No. PCT/GB96/01794.

Thus, in a first aspect the present invention provides a method for determining coat colour genotype in a pig which comprises:

- (a) obtaining a sample of pig nucleic acid; and
- (b) analysing the nucleic acid obtained in (a) to determine whether a mutation is/is not present at one or more exon/intron splice sites of the *KIT* gene.

In particular, the method determines whether a mutation is/is not present at the exon 17/intron 17 boundary, eg the substitution of the G of the conserved GT pair for A.

Reverse Transcriptase based Polymerase Chain analysis (RT-PCR) of the exon 16-19 region of *KIT* mRNA in animals of the Large White breed (*II/I*) and comparison to that transcribed in Hampshire animals (*i/i*) revealed an extra species of molecule in the former animals. RT-PCR analysis of both breeds yielded a product fragment with a length of 424bp indicating that both types of animal contained a *KIT* mRNA transcript containing a region corresponding to this region of the gene. However in addition the *II/I* animals yielded a RT-PCR product of 301bp indicating the presence of an mRNA species which did not contain the full transcription of the exon 16-19 DNA sequence (see example 1). These two transcripts have been shown to be derived from the separate duplicate copies of the *KIT* gene associated with the *I* allele. Sequencing over the *KIT* exon 17/intron 17 boundary revealed a difference in the intron boundary sequences present in the two duplicate copies of the *KIT* gene associated with the *I* allele (see example 2). The sequences are as shown below:

Allele	Kit Gene	Exon17 Intron 17
<i>i</i>	KIT1AAC GTG.....
<i>P</i>	KIT1AAC GTG.....
<i>P</i>	KIT2AAC GTG.....
<i>I</i>	KIT1AAC GTG.....
<i>I</i>	KIT2AAC ATG.....

This alteration of the 5' intron splice site from a GT pair to an AT pair affects the splicing of the pre mRNA and results in the loss of the whole of exon 17 from the mRNA transcribed from the *I*-KIT2 sequence. This will result in a modified KIT protein with the associated alterations in function and phenotype.

Based upon the sequence polymorphism rapid tests can be developed to determine the alleles carried by a specific animal at the dominant white locus.

- 5 In one form such a test would comprise amplification of the region through the polymerase chain reaction (PCR) utilising genomic DNA from the animal in question as a template. The nucleotide sequence CATG comprises a recognition sequence for the restriction enzyme *Nla*III. This sequence is only present in the *I-KIT* 2 sequence at the junction position and thus one can differentiate the
- 10 DNA molecules amplified from the two alleles by digestion of the amplification products with this enzyme or any other restriction enzyme with a suitable recognition site.

- Genomic DNA for use in such a test can be prepared by a wide range of
- 15 available methods, from any tissues or products derived from the animal in question. A 175bp fragment of the *KIT* gene containing the exon 17/intron 17 boundary region can be amplified from porcine genomic DNA using a pair of primers such as:

- 20 KIT21 (5'-GTA TTC ACA GAG ACT TGG CGG C-3'); and
KIT35 (5'-AAA CCT GCA AGG AAA ATC CTT CAC GG-3').

- The use of such primers in a PCR-RFLP test yields a fragment of 175bp before digestion with *Nla*III. In alleles with the G present at nucleotide position 1 of
- 25 intron 17 (*I-KIT*1 type sequence) there is only one *Nla*III cleavage site present

41bp from the end of the fragment. This site is present in all versions of the *KIT* sequence. Thus digestion of the 175bp fragments obtained from *i* and *P* alleles yields two fragments of 134bp and 41bp. Where the G is mutated to A as in the *I-KIT2* sequence present in *I* and *I** a further *Nla*III cleavage site is created and thus digestion yields products of 80bp and 54bp and 41bp (see figure 3, example 2). A number of other oligonucleotides suitable as PCR primers could easily be derived from the sequence of this region of the porcine genome. An example of such a PCR-RFLP test is given in example 3. The results that would be obtained from such a PCR-RFLP test described above are as shown below:

	Genotype	Fragment sizes	Fragment sizes
		134 + 41 bp	80 + 54 + 41 bp
	<i>III</i>	Yes	Yes
	<i>I/P</i>	Yes	Yes
15	<i>Ii</i>	Yes	Yes
	<i>P/P</i>	Yes	No
	<i>P/i</i>	Yes	No
	<i>ii</i>	Yes	No
	<i>I*/I</i>	Yes	Yes
20	<i>I*/I*</i>	Yes	Yes
	<i>i/I*</i>	Yes	Yes
	<i>I*/P</i>	Yes	Yes

Thus, simply by analysing restriction products certain genotypes can be distinguished. However, there are others which cannot be distinguished in this first configuration of the test. In a further refinement the test can be carried out in such a way that the amount of each fragment can be calculated. By carrying

out electrophoresis on an apparatus that allows the quantification of each of the bands one can determine the ratio of the two forms of template in the genomic DNA sample used. Examples of such an apparatus include the Perkin Elmer Applied Biosystems 373 and 377 DNA sequencing systems. The application of this type of equipment is illustrated in example 4. Any equipment capable of determining the relative amounts of the products from the two different sequences is equally applicable to such tests. The expected results from such a test are shown below.

Genotype	Normal KIT sequence Copies	Splice mutant KIT Copies	Ratio <u>Normal</u> Splice mutant
<i>III</i>	2	2	1
<i>I/P</i>	3	1	3
<i>II/I*</i>	2	2	1
<i>II/i</i>	2	1	2
<i>P/P</i>	4	0	0
<i>P/I*</i>	3	1	3
<i>P/i</i>	3	0	0
<i>I*/i</i>	2	1	2
<i>I*/I*</i>	2	2	1
<i>i/i</i>	2	0	0

10

Thus, using such a test one could identify all animals carrying alleles of dominant white that might dispose themselves or their offspring to exhibiting non-white coat colour (*i* or *P*) as those giving a ratio other than one. Depending on the requirement and the derivation of the lines under selection one could take

an appropriate subset of animals. For example in a cross derived animals carrying only alleles *I* and *i* one could identify any white individuals (*III* or *I/i*) carrying *i* as they would have a ratio in the test of 2 as opposed to 1 for the *III* animals. The distinction of alleles *I* and *I** can be carried out on the basis of the
5 4bp deletion as described in the previously filed patent publication WO97/05278.

There are a range of techniques by which differentiation of alleles containing the splice mutation and those containing the normal sequence could be
10 differentiated by a person expert in the field.

Analysis of the genetic composition of an animal could be based upon a number of different source materials. These include genomic DNA, RNA and the KIT protein itself. There may also be effects on the levels and nature of other
15 proteins, metabolites and RNA species which could be measured to create a more indirect assay.

DNA could be used as the basis for a number of approaches to testing. One approach is through the amplification of the region of DNA containing the
20 polymorphism using the polymerase chain reaction. This could then be linked to a number of forms of analysis of the product. Examples of electrophoresis based technologies include Single Strand Conformation Polymorphism (SSCP), Restriction Fragment Length Polymorphism (RFLP) and DNA sequencing of PCR products or direct genome sequencing. Other PCR based techniques that
25 might alternatively be used include the Perkin Elmer TaqMan systems, Single Nucleotide Polymorphic Extension (SNUPE) and Minisequencing. PCR products from the region might also have application in hybridization based

approaches to the differentiation of alleles at this locus. Hybridization methods might include the probing of Southern transfers of genomic DNA with allele specific oligonucleotide, RNA, DNA fragment or Protein Nucleic Acid (PNA) probes. Other strategies of application here include hybridization of genomic
5 DNA or PCR products (specific or whole genome) to oligonucleotide arrays possibly in the form of 'DNA chips'. Such arrays could consist of any reagent capable of binding DNA or RNA derived from the locus in question in an allele specific manner. Further useful methods of analysis also include oligonucleotide ligation assay and the ligase chain reaction. For a review on
10 methods for detecting point mutations see Landegren, 1996, Laboratory Protocols for Mutation detection, Oxford University press, Oxford.

A number of effects on the RNA produced from the gene in question have already and may in the future be observed. All such differences between the
15 mutated and normal forms of the gene are useful targets for the determination of genotype and a large range of methods are available to the person skilled in the art. The changes that are or might be observed and methods of analysis are as follows. Alteration of the size, rate of processing, stability and quantity of RNA transcripts could be measured through widely used techniques such as northern
20 blotting and RT-PCR as well as a number of the techniques described above for DNA analysis such as hybridization to oligonucleotide or DNA fragment arrays.

Another approach which can be used is to use a linked genetic polymorphism
25 which is closely associated with the presence or absence of the alteration at the exon/intron boundary. Such a polymorphism may occur in the *KIT* gene itself or in a chromosomal region linked to *KIT*. By using a single linked marker in

complete association with the presence/absence of the duplication or a combination of markers showing a partial association a highly informative test can be developed. For instance, the SSCP (Single Strand Conformation Polymorphism) method may be used to develop such polymorphism. The principle of the method is that double-stranded DNA, produced by PCR, is denatured into single-stranded DNA which is then separated by non-denaturing gel electrophoresis. Under non-denaturing conditions the single-stranded DNA forms a secondary structure due to intra-strand interaction but a proportion of the single-stranded DNA will renature and form double-stranded DNA. Two types of polymorphism may be revealed by this method. Firstly, a difference in nucleotide sequence between two alleles may influence the secondary structure of single-stranded DNA which is revealed as a difference in the mobility rate during electrophoresis. Secondly, a difference in nucleotide sequence often influences the mobility of the heteroduplex DNA (A heteroduplex is a double-stranded DNA molecule formed by two single-stranded molecules representing different alleles).

Association between genetic markers and genes responsible for a particular trait can be disrupted by genetic recombination. Thus, the closer the physical distance between the marker and the gene in question, the less likely it is that recombination will separate them.

It is also possible to establish linkage between specific alleles of alternative DNA markers and alleles of DNA markers known to be associated with a particular gene (e.g. the *KIT* gene discussed herein), which have previously been shown to be associated with a particular trait. Thus, in the present situation, taking the *KIT* gene, it would be possible, at least in the short term, to select for pigs with a particular coat colour, indirectly, by selecting for certain alleles of a *KIT* gene

associated marker through the selection of specific alleles of alternative chromosome 8 markers. Examples of such markers known to be linked to the *KIT* gene on porcine chromosome 8 include genetic polymorphism in the *KIT* gene itself or in the closely linked genes for the α -subunit of platelet derived growth factor (PDGFRA) and albumin.

Particular genetic markers associated with the *KIT* gene are microsatellites. These are simple sequence repeats of 4, 3 or, more usually, 2 nucleotides, which occur essentially at random around the genome at approximately every 50,000 bases (about 60,000 microsatellites per haploid genome). Stuttering of DNA polymerase during replication and unequal crossing-over during recombination are thought to result in the loss or gain of repeat units. This means that microsatellites are usually polymorphic and can have several repeat length alleles.

Examples of linked microsatellite sequences include S0086 (Ellegren *et al*, *Genomics*, 16:431-439 (1993)) , S0017 (Coppieters *et al*, *Animal Genetics* 24: 163-170 (1993)), Sw527, Swr750 and SW916 (Rhorer *et al*, *Genetics*, 136:231-245 (1994)) . It would be possible to select indirectly for alleles of the *KIT* gene linked to coat colour using any of the above markers, or indeed any other linked markers on porcine chromosome 8.

Alterations in the level of the KIT protein could be measured using either specific antibodies for example in an ELISA system or on western blots or through the use of a range of biochemical techniques to measure the activity of the protein. Such tests could also be applied to other proteins or metabolites, the level or nature of which is altered by the presence of specific alleles at the locus. Different protein structures due to the presence of specific alleles could

be identified through the use of structure specific antibodies. As with the DNA and RNA based methods all these protein methods could be applied in a quantitative fashion thus achieving the full discriminatory capabilities possible.

5 Kits could be produced for the specific analysis of the polymorphism described here alone and also with reagents allowing the combined analysis of the other polymorphisms previously reported and the subject of patent publication WO97/05278.

10 The invention will now be described with reference to the following examples, which should not be construed as in any way limiting the invention. The examples refer to the figures in which:

15 **Figure 1:** is a schematic representation of the structure of the known porcine *KIT* alleles, where 4bp del'n refers to the 4bp deletion in intron 18 of one copy of the duplicated *KIT* gene DNA as reported by Moller *et al.* 1996 and Exon 17-A refers to the change of nucleotide 1 of intron 17 from a G as in the wild type allele to an A as reported in this patent;

20 **Figure 2:** shows an electropherogram (4% agarose Nusieve/Seakem 3:1; 100V for 80 min) showing RT-PCR products of *KIT* exon 16-19 with the primers KIT1F and KIT7R. The samples 1-3 and 4-6 are Swedish Large White and Hampshire pigs, respectively. The size difference between the 424 and 301 bp fragments is due to lack of exon 17 in the latter
25 fraction. The two upper bands of the Yorkshire pigs were interpreted as heteroduplexes (HD);

Figure 3: shows a 48 bp sequence comprising 21 bp of KIT exon 17 and 27 bp of KIT intron 17 where the position of the intron/exon border is marked with a vertical line, the splice site mutation (nt1 $G \rightarrow A$) indicated with a vertical arrow and identical bases in alleles P and i are marked with a dot;

Figure 4: shows the results of *Nla*III PCR RFLP test used to detect the presence of a splice site mutation in intron 17 of the KIT gene. Figure 4A shows the position of two *Nla*III recognition sites within the PCR product amplified using primer pair KIT21 and KIT35. All distances are given in base pairs. Figure 4B shows the size of fragments which result following *Nla*III digestion of either normal *KIT* or splice mutant *KIT*. Figure 4C illustrates use of the PCR RFLP test. Lane 1 shows the KIT21/KIT35 amplified fragment undigested. Digestion was performed on PCR products amplified from, in Lane 2: a clone which contains the splice site mutation; Lane 3: a clone which contains the normal splice site sequence; Lane 4: genomic DNA from a coloured pig; Lane 5: genomic DNA from a white pig. Fragment sizes are given in base pairs;

Figure 5: shows a comparison of the ratio of normal to splice mutant *KIT* in animals of genotypes II , I/i and I/P ;

Figure 6: shows the ratio values for 56 Landrace and 33 Large White animals. A clearly bimodal distribution is observed with 7 Landrace and 3 Large White individuals having a ratio value of approximately 3 or above, suggesting them to be heterozygous carriers for the P allele (genotype I/P). This means P has gene frequency estimates of 6.25%

(7/112 chromosomes tested) and 4.5% (3/66 chromosomes tested) within the Landrace and Large White breeds respectively; and

FIGURE 7: shows a plot of Ct FAM versus Ct TET for animals of genotypes *Ii* and *II* analysed for KIT splice mutant genotype using TaqMan® chemistry.

Example 1

RT-PCR of porcine *KIT* exon 16-19

i. mRNA purification from blood samples

Fresh blood samples were collected in citrate tubes from coloured Hampshire pigs and Large White pigs. Leukocytes were isolated from 5 ml blood using Ficoll 100 (Pharmacia Biotech). Isolation of mRNA from leukocytes was then carried out using the Quickprep *Micro* mRNA purification kit (Pharmacia Biotech). The mRNA was stored as a precipitate under ethanol at -70°C for up to one month before use in reverse transcriptase (RT)-PCR.

ii. RT-PCR of KIT exon 16-19

First-strand cDNA synthesis was accomplished using the First-Strand cDNA Synthesis kit (Pharmacia Biotech) so that ~100 ng mRNA was randomly primed by 0.1 µg pd(N6) in a total volume of 15 µl. Two µl of the completed first cDNA strand reaction was then directly used per 12 µl PCR reaction by adding 10 µl PCR mix containing 10 pmol each of the mouse/human derived primers KIT1F and KIT7R (5'-TCR TAC ATA GAA AGA GAY GTG ACT C

and 5'-AGC CTT CCT TGA TCA TCT TGT AG, respectively; Moller et al. 1996, *supra*), 1.2 µl 10 x PCR-buffer (10 mM Tris-HCl, pH 8.3; 50 mM KCl) and 0.5 U of AmpliTaq polymerase (Perkin-Elmer) incubated with an equal amount Taqstart antibody (Clontech) at 25 °C for 5 min to achieve a hot start
5 PCR. The reaction was covered with 20 µl mineral oil and thermocycled in a Hybaid Touchdown machine (Hybaid) with 40 cycles at 94°C for 1 min, 55-48 °C (touchdown one degree per cycle the first seven cycles and then 48°C in the remaining cycles) for 1 min and 72°C for 1 min. After PCR 2µl loading dye was added to each sample which were then loaded on 4% agarose gel
10 (Nusieve/Seakem 3:1, FMC Bioproducts) and electrophoresed with 100V for 80 min. Products were visualised by ethidium bromide staining and UV-illumination.

iii. Cloning and sequencing of RT-PCR-products

15 The RT-PCR products representing *KIT* exon 16-19 were purified by extraction from 2% agarose gels using the QIAEX gel extraction kit (QIAGEN) and cloned into the pUC18 vector using the Sureclone ligation kit (Pharmacia Biotech). Plasmids were isolated using the QIAfilter plasmid Midi kit (QIAGEN). Cloned plasmid inserts were sequenced using dye primer
20 chemistry. Each cycling reaction was prepared with plasmid template DNA and ready reaction mix containing fluorescently labelled M13 forward or reverse primer as described in the ABI Prism protocol P/N 402113 (Perkin Elmer). Cycling and sample pooling were performed using a Catalyst 800 Molecular Biology Workstation (ABI) following the instruments user manual (Document
25 number 903877, Perkin Elmer). The resulting extension products were purified, loaded and analysed using the 377 ABI Prism sequencer as described by the instrument protocol P/N 402078 (Perkin Elmer).

iv Results and discussion

A 424 bp fragment including KIT cDNA exon 16-19 was amplified from all pigs. The Hampshire pigs did not show any additional products whereas the
5 Large White pigs (eight tested) all showed a 301 bp truncated cDNA fragment (Fig 2). Sequence analysis revealed the 424 bp fragment was identical in the two breeds whereas the whole exon 17 (123 bp) was missing from the 301 bp fragment. Apparent differences between individuals regarding the relative
10 amounts of these two products may have been caused either by different genotypes containing differing numbers of copies of the *KIT* gene sequence, individual differences in mRNA expression levels or random RT-PCR effects.

The two upper fragments present in Large white pigs represent heteroduplexes between the 301 and 424 bp fragments (Fig. 2). This was shown by an
15 experiment where these slow migrating fragments were generated by pooling homoduplexes of the 424 and 301 bp which were then heat denatured and cooled to 25°C. Moreover, cloning of the lower heteroduplex fraction of a Large White pig resulted in clones with insert length corresponding to either of the
20 two homoduplexes.

Example 2

PCR Amplification and Sequencing of KIT Exon 17-Intron 17 (5' Splice Site)

i. PCR to produce DNA Sequencing Template

25 A 175 bp region including the boundary between exon17 and intron17 of the *KIT* gene was amplified for sequence analysis using forward primer KIT21 (5' - GTA TTC ACA GAG ACT TGG CGG C -3') and reverse primer KIT35 (5' -

AAA CCT GCA AGG AAA ATC CTT CAC GG - 3'). PCR was carried out on a DNA thermal cycler (Perkin Elmer 9600) in a total volume of 20 μ l containing 25 ng genomic DNA, 1.0 mM MgCl₂, 50 mM KCl, 10 mM Tris-HCl, pH 8.3, 200 μ M dNTPs, 0.5 U AmpliTaq Gold (Perkin Elmer) and 10 pmol of both KIT21 and KIT35 primer. To activate AmpliTaq Gold, initial heat denaturation was carried out at 94°C for 10 minutes followed by 32 cycles each consisting of 45 sec at 94°C, 45 sec at 55°C and 45 sec at 72°C. The final extension lasted for 7 min at 72°C. PCR products were cloned into vector pUC18 using the SureClone ligation kit (Pharmacia Biotech).

ii. Preparation of Plasmid DNA

Plasmid DNA was purified from overnight bacterial culture using the Jetstar plasmid midi kit (Genomed) and the resulting DNA diluted to 150 ng/ μ l.

iii. Sequencing of plasmid DNA

DNA was sequenced as in example 1 section iii.

iv. Results

A portion of the DNA sequence from exon 17 and intron 17 of the KIT gene was determined and compared between animals with each of these three alleles. Figure 3 shows that the *I* allele carries a splice site mutation at position 1 of intron 17. This G to A base substitution is present in one of the two gene copies carried on each chromosome. The base substitution occurs in the invariant GT dinucleotide which characterises 5' exon/intron boundaries. Analysis of the *I*^P allele showed the splice site mutation was not present in either the normal (KIT1) or duplicated copy of the gene (KIT2). We have found the splice site

mutation is unique to the *I* alleles, and therefore makes it possible to distinguish the I-KIT2 sequences.

Example 3

5

Testing For the Presence of the Splice Site Mutation with PCR RFLP

To easily test for the presence of the G to A splice site mutation, restriction endonuclease *Nla*III (CATG) was used to exploit the point substitution identified at position 1 of intron 17 (Figure 3). The *Nla*III recognition sites in the fragment amplified from *KIT* and the expected restriction products are illustrated in Figure 4A and 4B respectively.

15

i. DNA preparation for RFLP Test

DNA can be prepared from any source of tissue containing cell nuclei, for example white blood cells, hair follicles, ear notches and muscle. The procedure here relates to blood cell preparations; other tissues can be processed similarly by directly suspending material in K buffer and then proceeding from the same stage of the blood procedure. The method outlined here produces a cell lysate containing crude DNA which is suitable for PCR amplification. However, any method for preparing purified, or crude, DNA should be equally effective.

25

Blood was collected in 50mM EDTA pH 8.0 to prevent coagulation. 50µl of blood was dispensed into a small microcentrifuge tube (0.5ml Eppendorf or

equivalent). 450µl of TE buffer was added to lyse the red blood cells (haem groups inhibit PCR) and the mix vortexed for 2 seconds. The intact white and residual red blood cells were then centrifuged for 12 seconds at 13,000 g in a microcentrifuge. The supernatant was removed by gentle aspiration using a low pressure vacuum pump system. A further 450µl of TE buffer was then added to lyse the remaining red blood cells and the white blood cells collected by centrifugation as before. If any redness remained in the pellet, this process was repeated until the pellet was white. After removal of the last drop of supernatant from the pelleted white blood cells, 100µl of K buffer containing proteinase K was added and the mixture incubated at 55 degrees C for 2 hours. The mixture was then heated to 95-100 degrees C for 8 minutes and the DNA lysates stored at -20°C until needed.

Reagents

15 T.E. Buffer: 10mM TRIS-HCl pH8.0
1mM EDTA

K Buffer: 50mM KCl
10mM TRIS-HCl pH8.3
20 2.5mM MgCl₂
0.5% Tween 20

ii. Restriction Enzyme Digestion and Electrophoresis

The PCR amplification product is 175 bp in length. To test for polymorphism at position 1 of intron 17, digestion reactions were set up as below:

3.0 µl PCR amplified DNA

1.0 µl 10 X NEBuffer 4
0.1 µl BSA 100 µg/ml
0.1 µl *Nla*III 10 U/µl
5.8 µl dH₂O

5

(1 X NEBuffer 4 (New England Biolabs) contains 50 mM potassium acetate, 20 mM Tris acetate, 10 mM magnesium acetate and 1 mM DTT). Following incubation at 37°C for 90 minutes each 10 µl reaction volume had 2 µl of loading dye added and the mix loaded on a 8% native polyacrylamide gel (Protagel, 37.5:1 acrylamide:bisacrylamide, National Diagnostics, Atlanta) in 10. 0.5 X TBE (44.5 mM Tris pH 8.0, 44.5 mM boric acid and 0.5 mM EDTA) and electrophoresed for 3 hours at 200V in a vertical slab unit (SE600 Hoefer Scientific Instruments). Products were visualised by ethidium bromide staining.

15 iii Results

A PCR RFLP protocol was designed to test for the presence of the splice site mutation as the substitution occurs within the recognition site for restriction endonuclease *Nla*III. Figure 4B illustrates that presence of the G to A base substitution at position 1 of KIT intron 17 results in restriction at each of two 20 *Nla*III recognition sites within the 175 bp DNA fragment. Following electrophoresis, this results in fragments of sizes 80 bp, 54 bp and 41 bp. Where the splice site mutation is absent however, incubation with *Nla*III results in digestion only at recognition site 1. Following electrophoresis this results in fragments of 134 bp and 41 bp. The invariant *Nla*III recognition site 1 serves as 25 an internal control to ensure complete digestion has taken place. Results of this PCR RFLP analysis are illustrated in Figure 4C. Analysis was performed on fragments amplified from clones which either carry the splice site mutation (lane 2) or carry the normal splice site sequence (lane 3). Lane 4 shows the result of analysis where DNA amplified from the genomic DNA of a coloured

animal was used. Lane 5 shows the resulting bands where a white animal was tested. The test was used to analyse 121 individuals from seven different breeds of pig. The splice site mutation was found only in the 97 animals with the dominant white phenotype (*I*/- or *I**/*i*) and none of the 24 coloured (*I*^P or *i*) examples (Table 2). This analysis confirms *I* and *I** to be unique in that they are the only alleles to carry the splice site mutation.

TABLE 2

Distribution of the Splice Site Mutation Between Different Breeds and Coat Phenotype

Breed	Coat Colour	Assumed Genotype ¹	Animals Tested	Normally spliced KIT ²	Splice Mutation ²
Large White	White	<i>I</i> /-	33	33	33
Landrace	White	<i>I</i> /-	56	56	56
Hampshire	Coloured	<i>i</i> / <i>i</i>	5	5	0
Duroc	Coloured	<i>i</i> / <i>i</i>	5	5	0
Pietrain	Coloured	<i>i</i> / <i>i</i>	8	8	0
Meishan	Coloured	<i>i</i> / <i>i</i>	5	5	0
Wild Boar	Coloured	<i>i</i> / <i>i</i>	1	1	0
Wild Boar	White	<i>I</i> */-	8	8	8
x Large White					
Totals					
	White	<i>I</i> /-	89	89	89
	White	<i>I</i> */-	8	8	8
	Coloured	<i>i</i> / <i>i</i>	24	24	0

¹ White animals may be homozygous or heterozygous for the *I* allele

² Presence of the splice site mutation determined by *Nla*III PCR RFLP test

Example 4**Quantification of Normal KIT and Splice Mutant KIT (Intron 17 nt1^{G→A})**

- 5 As the splice site mutation is present in only one of the duplicated regions of *I* and not in the duplicated region of *I^P*, the various genotypes can be expected to have the attributes described in Table 3.

TABLE 3

Genotype	Copies of Normal KIT	Copies of KIT containing the splice mutation	Ratio of normal KIT to splice mutant KIT
<i>I/I</i>	2	2	1:1
<i>I/i</i>	2	1	2:1
<i>i/i</i>	2	0	2:0
<i>I/I^P</i>	3	1	3:1
<i>I^P /i</i>	3	0	3:0

10

- Due to the dominance of allele *I*, three of the genotypes in Table 2 are carried by white animals and therefore can not be identified by phenotypic characterisation. Quantification of the relative amounts of the normal *KIT* gene and the splice mutant *KIT* gene allows the ratio between the two to be calculated, and therefore the genotype of individual animals predicted. This was achieved by quantification of two DNA fragments following *Nla*III digestion.
- 15

The amount of 134 bp fragment, representative of the normally spliced *KIT* gene, and of 54 bp fragment, representative of the splice mutant *KIT*, were measured following electrophoresis using GeneScan software.

5 i. PCR to Produce DNA for Quantification

As described in example 2 section i. The reverse primer KIT35 is labelled with the ABI fluorescent dye FAM at the 5' end.

ii Restriction Enzyme Digestion

10 As described in example 2 section ii.

iii Electrophoresis and Quantification of DNA Fragments

Following digestion, 0.5 µl of the reaction volume was mixed with 2.5 µl of deionised formamide, 0.5 µl of GS350 DNA standard (ABI) and 0.4 µl blue
15 dextran solution before being heated to 90°C for 2 minutes and rapidly cooled on ice. Three µl of this mix was then loaded onto a 377 ABI Prism sequencer and the DNA fragments separated on a 6% polyacrylamide gel in 1 X TBE buffer for 2 hours at 700 V, 40 mA, 32 W. The peak area of fragments representative to both the normal and splice mutant forms of *KIT* were
20 quantitated using the GeneScan (ABI) software.

iv. Ratio Calculations

The peak area value of the 134 bp fragment (normal *KIT*) was divided by twice the peak area value of the 54 bp fragment (splice mutant *KIT*) in order to
25 calculate the ratio value for each sample.

v. Results

Analysis was performed on animals from the Swedish wild pig/Large White intercross pedigree for which genotypes at *I* have been determined by conventional breeding experiments with linked markers. Figure 5 and Table 4 show the ratio of normal to mutant *KIT* calculated for animals from each of the three genotype classes, *I/I* (expected ratio 1:1), *I/i* (expected ratio 2:1) and *I/I^P* (expected ratio 3:1). The results are entirely consistent with the expected ratio values and indicate that the three genotype classes can be distinguished using this method.

TABLE 4

Ratio of the Two *KIT* Forms in Different Dominant White Genotypes in a Wild Pig/Large White Intercross

Genotype	Phenotype	Expected Ratio (Normal: Mutant)	Observed Ratio (Normal: Mutant) ± SE	Number Tested
<i>I/I</i>	White	1:1	1.15 ± 0.075	13
<i>I/I^P</i>	White	3:1	3.11 ± 0.084	12
<i>I/i</i>	White	2:1	2.23 ± 0.109	14

Figure 5 illustrates that the range of ratio values calculated for the two genotypes *I/I* and *I/I^P* do not overlap. This enables animals carrying the *I^P* allele to be identified and the frequency of the allele within different pig breeds determined. Ratio values were calculated for 56 Landrace and 33 Large White animals and the results are shown in Figure 6. A clearly bimodal distribution is

observed with 7 Landrace and 3 Large White individuals having a ratio value of approximately 3 or above, suggesting them to be heterozygous carriers for the I^P allele (genotype I/I^P). This means I^P has gene frequency estimates of 6.25% (7/112 chromosomes tested) and 4.5% (3/66 chromosomes tested) within the Landrace and Large White breeds respectively.

Example 5

Analysis for presence and quantification of the porcine KIT splice mutation using the PE ABI TaqMan chemistry

Method

i. Preparation of template DNA for PCR

DNA was prepared as in example 3, section i

ii. TaqMan[®] PCR reactions

TaqMan[®] PCR reactions were set up as shown in table 5

TABLE 5

PCR mix for TaqMan[®] based splice mutation test

Reagent	Final Conc ⁿ	Volume
10x TaqMan [®] Buffer A (Perkin Elmer)	1x	2.50 µl
25mM MgCl ² Sol ⁿ	5mM	5.00 µl
DATP	200µM	0.50 µl
DCTP	200µM	0.50 µl
DGTP	200µM	0.50 µl
DUTP	200µM	0.50 µl
Amplitaq Gold TM (5U/µl) (Perkin Elmer)	0.05U/µl	0.25 µl
AmpErase TM N-Glycosylase (1U/µl) (Perkin Elmer)	0.01U/µl	0.25 µl
KITTM -NEST-F (5µM)	500nM	2.50 µl
KITTM-NEST-R (5µM)	500nM	2.50 µl
KITTM FAM (5µM)	100nM	0.50 µl
KITTM TET (5µM)	100nM	0.50 µl

25% Glycerol	8%	8.00 μ l
Porcine genomic DNA		<u>1.00 μl</u>
		25.00
		μ l

The PCR primers used were as described below:

KITTM-Nest-F (5'-CTC CTT ACT CAT GGT CGA ATC ACA-3') and

KITTM -Nest-R (5'-CGG CTA AAA TGC ATG GTA TGG-3').

The TaqMan[®] probes used were:

KITTM-A FAM (5'-TCA AAG GAA ACA TGA GTA CCC ACG CTC-3') and

KITTM -G TET (5'- TCA AAG GAA ACG TGA GTA CCC ACG C -3')

The TaqMan[®] probes were prepared by Perkin Elmer and labelled with FAM and TET as indicated as well as the standard quenching group TAMRA. The 10x TaqMan[®] Buffer A, Amplitaq Gold[™], AmpErase N-Glycosylase, NTP's and 25mM MgCl₂ used were part of the TaqMan[®] PCR Core reagent Kit, supplied by Perkin-Elmer.

The reactions were then placed into a Perkin Elmer ABI Prism 7700 Sequence Detector and the reaction carried out using the following thermal profile, 50°C for 2 minutes, 95°C for 10 minutes followed by 40 cycles of 95°C 15s, 62°C 60s. The reactions were carried out under the control of 'Sequence Detector V.1.6' software using the 'Single Reporter' and Real-Time' options with the 'Spectral Compensation' function activated. Upon completion of the run real-time profiles for each sample were examined on the ABI7700 to check for any samples giving highly irregular profiles which were then excluded. The thresholds for both dyes, Fam and Tet, were set so that they intercepted each

dye during the exponential phase of PCR. Following updating of the calculations in 'Sequence Detector V.1.6' software results were exportated into MS Excel for further analysis.

5 iii. Analysis of results

Based upon the underlying theoretical principle that one cycle of PCR gives a doubling in the amount of cleavage of the quenching dye from the allele specific probe and therefore doubles the signal one would expect the threshold
10 cycle numbers from the *II* and *Ii* genotypes analysed to be as below:

Table 6:

15 Theoretical results for TaqMan® analysis of genotype at the KIT splice mutation

Genotype	Copies KIT 1 (G)	Copies KIT 2 (A)	Theoretical Ct TET (G)	Theoretical Ct FAM (A)
<i>II</i>	2	2	X	Y
<i>Ii</i>	2	1	X	Y+1

In theory the Ct for TET and FAM signals, represented as X and Y should be the same, as equal numbers of copies of the target sequences should be present in an *II* animal. However in practice this does not necessarily occur due to
20 differences in the hybridization and cleavage efficiency of the two probes and variation in the setting of the threshold cycle between the two dye signals. The reduction in splice mutant containing (A) sequences relative to those not containing the splice mutation (G) in the *Ii* animals ie 2:1 G:A ratio rather than

1:1 as for *II* genotype, should lead to the FAM signal reaching the threshold 1 cycle later than the TET signal in the genotype *Ii* animals. The actual results for samples tested are shown in Table 7.

5 **Table 7**
Ct values from analysis of *II* and *Ii* genotypes

Sample	Genotype	Ct FAM (A)	Ct TET (G)	Ct FAM - Ct TET
1	<i>Ii</i>	24.68	22.59	2.09
2	<i>Ii</i>	25.98	23.62	2.36
3	<i>Ii</i>	26.54	25.57	0.97
4	<i>Ii</i>	27.37	24.78	2.59
5	<i>Ii</i>	24.94	21.61	3.33
6	<i>Ii</i>	25.68	22.1	3.58
Ii Mean = 2.49				
7	<i>II</i>	22.05	23.78	-1.73
8	<i>II</i>	24.22	24.59	-0.37
9	<i>II</i>	24.19	23.85	0.34
10	<i>II</i>	23.66	23.51	0.15
11	<i>II</i>	24.35	22.71	1.64
12	<i>II</i>	22.82	21.69	1.13
13	<i>II</i>	22.84	22.7	0.14
14	<i>II</i>	23.17	22.9	0.27
Mean = 0.20				

No Template	35	35	0
No Template	35	35	0
No Template	35	35	0
No Template	35	35	0

Despite variation around the mean values it can be seen from Table 7 that there is a significantly increased delay in the FAM signal reaching the threshold level (approximately 2 cycles) relative to the TET signal in *Ii* animals compared to *II* animals as predicted, reflecting the reduced number of copies of the splice mutant (A) sequence present in animals of the *Ii* genotype. Plotting of the individual samples on a scatter plot (Figure 7) shows clustering of the two genotypes with the *Ii* cluster shifted along the Ct FAM axis due to the reduced number of copies of the KIT2 (A) sequence for which the FAM probe is specific.

CLAIMS:

1. A method for determining coat colour genotype in a pig which comprises:
 - 5 (a) obtaining a sample of pig nucleic acid; and
 - (b) analysing the nucleic acid obtained in (a) to determine whether a mutation is/is not present at one or more exon/intron splice sites of the *KIT* gene.
- 10 2. A method as claimed in claim 1 wherein the analysis in step (b) is carried out to determine whether a mutation is/is not present at the exon 17/intron 17 boundary.
3. A method as claimed in claim 2 wherein the mutation consists of the
15 substitution of the G of the conserved GT pair for A.
4. A method as claimed in any one of claims 1 to 3 wherein the sample of nucleic acid is amplified prior to analysis.
- 20 5. A method as claimed in claim 4 wherein the nucleic acid is genomic DNA.
6. A method as claimed in claim 5 wherein amplification is carried out using PCR and at least one pair of suitable primers.
- 25 7. A method as claimed in claim 6 wherein the pair of suitable primers is:

5'-GTA TTC ACA GAG ACT TGG CGG C-3'); and

5'-AAA CCT GCA AGG AAA ATC CTT CAC GG-3'.

8. A method as claimed in any one of claims 5 to 7 wherein after
5 amplification the nucleic acid is treated with a restriction enzyme, followed by
analysis of fragment lengths.

9. A method as claimed in claim 8 wherein the nucleic acid is treated with
the restriction enzyme *Not*III.

10

10. A method as claimed in claim 8 or claim 9 wherein the ratio of
restriction fragment lengths is determined.

11. A method as claimed in claim 4 wherein the nucleic acid is mRNA.

15

12. A method as claimed in claim 11 wherein the nucleic acid is amplified
using RT-PCR.

13. A method as claimed in claim 12 wherein the length of RT-PCR product
20 is determined.

14. A method for determining coat colour genotype in a pig which comprises the
step of analysing a sample of pig KIT protein to determine whether the protein is the
splice variant protein.

15. A kit for use in determining the coat colour genotype of a pig which comprises one or more reagents suitable for determining whether a mutation is present at one or more exon/intron splice sites of the *KIT* gene.

5

16. A kit as claimed in claim 15 which comprises one or more reagents for carrying out PCR and one or more pairs of suitable primers.

10

17. A kit as claimed in claim 16 which comprises the following pair of primers:

5'-GTA TTC ACA GAG ACT TGG CGG C-3'); and

5'-AAA CCT GCA AGG AAA ATC CTT CAC GG-3'.




FIG.1

2/8




FIG.2



3/8

Sequence Alignment Across the Exon /Intron Border of KIT Exon 17

Allele	Gene Copy	Sequence																
		Exon 17								Intron 17								
I	KIT1	AAT	TAC	GTG	GTC	AAA	GGA	AAC	↓	GTG	AGT	ACC	CAC	GCT	CTC	CTG	ACA	GTC
	KIT2	A
I ^p	KIT1	G
	KIT1	G
i	KIT1	G

FIG.3

4/8



5/8




FIG. 4(c)

6/8

Ratio of normal to splice mutant KIT




FIG.5

7/8


Ratios for Landrace and Large White

FIG.6

8 / 8

Plot of Ct FAM vs Ct TET for TaqMan based PCR
analysis of porcine KIT splice mutation genotype





FIG.7

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03081

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12Q1/68 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 05278 A (DALGETY PLC ;ANDERSSON LEIF (SE); MOLLER MARIA JOHANSSON (SE); WAL) 13 February 1997 cited in the application see the whole document ---	1,4-6
A	JOHANSSON MOLLER M ET AL: "Pigs with the dominant white color phenotype carry a duplication of the kit gene encoding the mast/stem cell growth factor receptor" MAMMALIAN GENOME, vol. 7, no. 11, November 1996, pages 822-30, XP002091653 see the whole document --- -/--	1,4-6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 January 1999

Date of mailing of the international search report

16/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Osborne, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03081

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>MARKLUND S ET AL: "Molecular basis for the dominant white phenotype in the domestic pig"</p> <p>GENOME RESEARCH,</p> <p>vol. 8, no. 8, 1998, pages 826-33,</p> <p>XP002091654</p> <p>see the whole document</p> <p>-----</p>	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03081

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9705278 A	13-02-1997	AU 6621296 A	26-02-1997
		CA 2227826 A	13-02-1997
		CN 1196755 A	21-10-1998
		EP 0842296 A	20-05-1998

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.